



Green and Highly Efficient Synthetic Approach for the Synthesis of 4-Aminoantipyrene Schiff Bases

Kovan Dilawer Issa ^{a*}, Rostam Rasul Braiem ^a

^aDepartment of Chemistry, College of Science, Salahaddin University, Erbil, Kurdistan Region-Iraq

ARTICLE INFO

Article history:

Received: 13 June 2022

Received in revised form: 27 June 2022

Accepted: 28 June 2022

Available online: 12 September 2022

Keywords:

4-Aminoantipyrene

Schiff Base

Pyrazolone

Organocatalyst

DMEA

ABSTRACT

The analog of 4-aminoantipyrene (4-amino-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazole-3-one), is interesting in the field of pharmacology and pharmaceutical chemistry and plays an important role as the ingredients for the production of the drugs and medicines. In the current study, a series of Schiff (including two new compounds) bases have been synthesized from 4-aminoantipyrene and substituted benzaldehydes via a simple and easy procedure in a short time with high yields. The structures of products have been characterized by (FTIR, ¹HNMR, ¹³CNMR, and Elemental Analysis).

1. Introduction

Heterocyclic compounds are major constituents of the drugs and treatments for several diseases, also they are used as antibacterial [1], antifungal [2], anticancer [3], antioxidant [4], anti-inflammatory [5], antipyretic [6], etc. Among the drug formulations, pyrazoles and their derivatives are interesting to be studied because of having a major role in the pharmaceutical and biologically active compounds [7]. Pyrazolones are considered typical derivatives of pyrazoles, in which their application, synthesis, and reactivity are reported. Their important biological activity and medicinal chemistry, such as antimicrobial [8], analgesic, antipyretic [9], antidepressant [10], antitumor [11], etc. as well as their vital characteristic in coordination chemistry [12], have been an important choice for the chemists [13].

The most famous pyrazolone derivative is (4-amino-1,5-dimethyl-2-phenyl-3-pyrazolone) which is known commonly as 4-aminoantipyrene or aminophenazone, a five-membered heterocyclic lactam ring with a lot of biological activities including anti-inflammatory [14], anthelmintic [15], antiviral [16], antibacterial [17], anticancer, and antifungal activity [18]. Among the chemists, new types of chemotherapy drugs that contain Schiff bases are significant consideration and particularly scavenger has been recognized against reactive oxygen and nitrogen species, as well as inhibiting oxidative burst of neutrophils [29]. Also, it has been utilized in

in the case of having aminoantipyrene in their structure to distinguish liver disease and clinical treatments [19].

In 1864, Hugo Schiff has synthesized new compounds containing azomethine group ($R_1-N=C-R_2R_3$) by the reaction of primary amines (R_1-NH_2) with aldehydes ($O=CHR_2$) or ketones ($O=C-R_2R_3$) in which R could be aliphatic or aromatic substituents, the product named after him and called (Schiff base) [20]. Schiff bases are famous to possess biological activities including antimicrobial, antiviral, anticancer, and anti-inflammatory activity [21, 22]. Also, their free radical scavenging effect has been reported by several researchers [23]. On one side, Schiff bases are also utilized as intermediates in organic synthesis [24], catalysts [25], polymer stabilizers, dyes, and pigments [26].

A wide number of Schiff bases can form metal complexes and have a vital role as biologically active compounds, this is a declaration that Schiff bases are not only influenced by organic and coordination chemistry, but they are present in the development of inorganic biochemistry, optical materials, and catalysis [27].

4-Aminoantipyrene can form Schiff bases with aldehydes directly [28] and their importance as free radical the protection of prophylactic of many illnesses and oxidative pressure including cancer [20].

* Corresponding author.; Tel.: +9647507809682; e-mail: kovan.dilawer@tiu.edu.iq

Organic catalysts (organocatalysts) are quite important due to several characteristics including availability, easy preparation, selectivity, working under mild conditions, and compatibility with many functional groups [30]. Therefore, an organocatalyst namely N,N-dimethyl ethanolamine (DMEA) (Figure 1) has been selected to produce the desired products, nevertheless there is no experiments for the synthesis of Schiff bases using the mentioned catalyst [31].




Fig. 1. Structure of N, N-Dimethylethanolamine (DMEA)

The aim of this work is utilization of an organic catalyst (DMEA) for the synthesis of some 4-aminoantipyrene Schiff bases by a simple, efficient, and easy method and in a short duration of time with excellent yield of the products, as well as testing their antioxidant activity.

2. Experimental

2.1. General

All the starting materials have been purchased from Sigma Aldrich, Merck, or Fluka, and used without further purification. Melting points were measured via (Electrothermal 9100) capillary melting point apparatus. Thin layer Chromatography (TLC) plate precoated with silica gel aluminum was used to detect the completion of the reaction with system eluent of (n-Hexane: Ethyl acetate 60:40), as well as utilizing the ultraviolet light at the wavelength of 354 nm for visualization. Infrared spectra have been recorded with KBr disc on FTIR (Schimadzu IR Affinity-1) Spectrophotometer. ¹H NMR and ¹³C NMR spectra were recorded in DMSO-*d*₆ on Varian Inova 500 MHz and 125 MHz, respectively. Elemental analysis was done on LECO model TruSpec CHN analyzer as shown in Electronic Supplementary Information (ESI).

2.2. Synthesis of Schiff Bases

Equimolar of 4-aminoantipyrene (1mmol) and substituted benzaldehydes (1mmol) was mixed in a round bottom flask and (1 mol %) of dimethyl ethanol amine (DMEA) has been added as a homogeneous catalyst. The mixture was stirred at room temperature for appropriate time as mentioned in table 1, the progress of the reaction has been monitored by TLC for disappearance of starting materials. After completion of the reaction the products were separated by addition of water (15 ml) and stirring for 10 minutes. After that the products were filtered out and finally, they are purified by recrystallization with ethanol.

All the products were characterized by melting point, FTIR, ¹HNMR, ¹³CNMR, and elemental analysis (C, H, N).

2.2.1. 4-(benzylideneamino)-1, 5 - dimethyl -2-phenyl-1, 2-dihydro-3H-pyrazol-3-one (C1)

Yield (93 %); White solid; m.p. 169–171 °C, IR_{vmax}cm⁻¹: 3037, 2941, 1654, 1595, 1568. ¹HNMR (500 MHz, DMSO, *d*₆) δ 2.46 (s, 3H, CH₃), 3.17 (s, 3H, CH₃), 7.39–7.83 (m, 10 H, Ar), 9.62 (s, H, CH), ¹³CNMR (DMSO, *d*₆, 125 MHz): δ 10.15 (CH₃), 35.82 (CH₃), 116.78 (C=C), 124.84, 127.69, 128.69, 129.62, 130.62, 135, 138.15, 152.70 (C=C), 154.80 (C=O), 160.08 (C=N). Anal. Calcd. for C₁₈H₁₇N₃O: C: 74.20, H: 5.88, N: 14.42 %. Found: C: 74.18, H: 5.90, N: 14.44 %.

2.2.2. 4-(4-fluorobenzylideneamino)-1,5-dimethyl-2-phenyl-1, 2-dihydro-3H-pyrazol-3-one (C2)

Yield (95 %); White crystals; m.p. 229–231 °C, IR_{vmax}cm⁻¹: 3070, 2941, 1651, 1598, 1568, 1502. ¹H-NMR (500 MHz, DMSO, *d*₆) δ 2.59 (s, 3H, CH₃), 3.31 (s, 3H, CH₃), 7.40–8.02 (m, 9 H, Ar), 9.7 (s, H, CH). ¹³C NMR (DMSO, *d*₆, 125 MHz): δ 9.72 (C, CH₃), 35.30 (C, CH₃), 115.64, 115.86, 124.58, 126.90, 129.13, 129.26, 129.34, 134.11, 134.14, 134.49, 152.16, 152.94 (=C-N), 159.51 (C=O), 162.00 (C=N), 164.46 (C-F). Anal. Calcd. for C₁₈H₁₆FN₃O: C: 69.89, H: 5.21, N: 13.58 %. Found: C: 69.88, H: 5.24, N: 13.57 %.

2.2.3. 4-(2-chlorobenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C3)

Yield (94 %); Pale yellow; m.p. 192–193 °C, IR_{vmax}cm⁻¹: 3049, 2941, 1649, 1581, 1566. ¹HNMR (500 MHz, DMSO, *d*₆) δ 3.17 (s, 3H, CH₃), 3.38 (s, 3H, CH₃), 7.39–7.83 (m, 10 H, Ar), 9.62 (s, H, CH), ¹³CNMR (DMSO, *d*₆, 125 MHz): δ 11.16 (CH₃), 36.36 (CH₃), 117.38 (=C-N) 126.19 (2C, Ar), 127.92 (1C, Ar), 128.40 (1C, Ar), 128.66 (1C, Ar), 130.42 (2C, Ar) 131.15 (1C, Ar), 132.63 (1C, =C-N), 135.67 (C=C-N), 150.17 (C-Cl), 151.07 (=C-N), 153.9 (C=O), 160.58 (C=N). Anal. Calcd. for C₁₈H₁₆ClN₃O: C: 66.36, H: 4.95, N: 12.90 %. Found: C: 66.35, H: 4.97, N: 12.88 %.

2.2.4. 4-(2,6-dichlorobenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C4)

Yield (95 %); Pale yellow solid; m.p. 196–197 °C, IR_{vmax}cm⁻¹: 3061, 2924, 1651, 1581, 1560. ¹HNMR (500 MHz, DMSO, *d*₆): δ 2.42 (s, 3H, CH₃), 3.22 (s, 3H, CH₃), 7.39–7.53 (m, 10 H, Ar), 9.77 (s, H, CH), ¹³CNMR (DMSO, *d*₆, 125 MHz): δ 10.85 (CH₃), 36.05 (CH₃), 110.2 (C=C), 126.26, 128.7, 130.43, 131.84, 134.69, 151.07 (C=C), 153.91 (C=O), 160.52 (C=N). Anal. Calcd. for C₁₈H₁₅Cl₂N₃O: C: 60.02, H: 4.20, N: 11.66 %. Found: C: 59.99, H: 4.22, N: 11.64 %.

2.2.5. 4-(4-hydroxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C5)

Yield (94 %); Brown crystals; m.p. 225–226 °C, IR_{vmax}cm⁻¹: 3057, 2924, 1604, 1577, 1558, 1512. ¹HNMR (500 MHz, DMSO, *d*₆) δ 2.40 (s, 3H, CH₃), 3.10 (s, 3H, CH₃), 6.83–7.64 (m, 9H, Ar), 9.48 (s, H, CH), 9.93 (s, H, OH). ¹³CNMR (DMSO, *d*₆, 125 MHz): δ 10.13 (CH₃), 36.05 (CH₃), 116.54 (C=C), 117.51,

124.84, 127.28, 129.72, 135, 152.17, 155.32 (C=O), 160.10 (C–OH), 160.39 (C=N). Anal. Calcd. for C₁₈H₁₇N₃O₂: C: 70.34, H: 5.58, N: 13.67 %. Found: C: 70.37, H: 5.56, N: 13.64 %.

2.2.6. 4-(3,4-dimethoxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C6)

Yield (95 %); Pale yellow; m.p. 185–187 °C, IR_{vmax}cm⁻¹: 3076, 2964, 1649, 1595, 1573, 1510. ¹HNMR (500 MHz, DMSO, d₆) δ 2.44 (s, 3H, CH₃), 3.13 (s, 3H, CH₃), 3.79 (s, 3H, CH₃), 3.82 (s, 3H, CH₃), 7.02 (s, 3H, CH₃), 7.27–7.53 (m, 10 H, Ar), 9.5 (d, H, CH), ¹³CNMR (DMSO, d₆, 125 MHz): δ 11.11 (CH₃), 36.8 (CH₃), 56.63, 56.82, 109.79, 112.7, 118.04, 123.23, 125.57, 127.95, 130.36, 131.79, 135.97, 150.29, 150.31, 152.16, 153.13, 155.82 (C=O), 161.11 (C=N). Anal. Calcd. for C₂₀H₂₁N₃O₃: C: 68.36, H: 6.02, N: 11.96 %. Found: C: 68.33, H: 6.03, N: 11.97%.

2.2.7. 4-(4-hexyloxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C7)

Yield (93 %); Pale yellow; m.p. 189–191 °C, IR_{vmax}cm⁻¹: 2953, 2941, 1637, 1608, 1598, 1568. ¹HNMR (500 MHz, DMSO, d₆) δ 3.17 (s, 3H, CH₃), 3.38 (s, 3H, CH₃), 7.39–7.83 (m, 10 H, Ar), 9.62 (s, H, CH), ¹³CNMR (DMSO, d₆, 125 MHz): δ 9.76 (CH₃), 13.90 (CH₃), 22.05, 25.15, 28.56, 30.97 (CH₂), 35.51 (CH₃), 67.57 (O–CH₂), 114.60 (=C–N), 116.76, 124.29, 126.68, 128.85, 129.09 (C, Ar), 130.12, 134.65 (=C–N), 151.84 (C=O), 154.30 (=C–O), 160.46 (C=N). Anal. Calcd. for C₂₄H₂₉N₃O₂: C: 73.63, H: 7.47, N: 10.73 %. Found: C: 73.60, H: 7.50, N: 10.70 %.

2.2.8. 4-(2-fluorobenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C8)

Yield (95 %); Pale yellow; m.p. 176–177 °C, IR_{vmax}cm⁻¹: 3059, 2941, 1651, 1595, 1573, 1517. ¹HNMR (500 MHz, DMSO, d₆) δ 3.19 (s, 3H, CH₃), 3.38 (s, 3H, CH₃), 7.39–7.83 (m, 10 H, Ar), 9.62 (s, H, CH), ¹³CNMR (DMSO, d₆, 125 MHz): δ 10.14 (CH₃), 35.59 (CH₃), 117.19 (C=C), 125.15, 125.3, 126.45, 127.52, 129.59, 132.36, 134.85, 146.84 (C=C–N), 152.78 (C=O), 159.83 (C=N). Anal. Calcd. for C₁₈H₁₆FN₃O: C: 69.89, H: 5.21, N: 13.58 %. Found: C: 69.91, H: 5.20, N: 13.56 %.

2.2.9. 4-(4-nitrobenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C9)


Yield (92 %); Yellow solid; m.p. 253–255 °C, IR_{vmax}cm⁻¹: 3051, 2932, 1647, 1575, 1554, 1519. ¹HNMR (500 MHz, DMSO, d₆) δ 3.23 (s, 3H, CH₃), 3.27 (s, 3H, CH₃), 7.36–7.55 (m, 10 H, Ar), 9.66 (s, H, CH), ¹³CNMR (DMSO, d₆, 125 MHz): δ 10.96 (CH₃), 35.82 (CH₃), 109.99 (C=C), 125.26, 126.5, 129.14, 130.48, 152.17, 153.91 (C=O). Anal. Calcd. for C₁₈H₁₆N₄O₃: C: 64.28, H: 4.79, N: 16.66 %. Found: C: 64.29, H: 4.76, N: 16.65%.

2.2.10. 4-(4-methoxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one (C10)

Yield (94 %); White solid; m.p. 181–182 °C, IR_{vmax}cm⁻¹: 3051, 2966, 1645, 1608, 1593, 1508. ¹HNMR (500 MHz, DMSO, d₆) δ 3.13 (s, 3H, CH₃), 3.79 (s, 3H, CH₃), 7.39–7.83 (m, 10 H, Ar), 9.62 (s, H, CH), ¹³CNMR (DMSO, d₆, 125 MHz): δ 10.23 (CH₃), 35.99 (CH₃), 55.75, 114.66, 117.28 (C=C), 124.80, 127.15, 129.31, 129.54, 130.81, 135.18, 152.29, 154.8, 160.30 (C=O), 161.49 (C=N). Anal. Calcd. for C₁₉H₁₉N₃O₂: C: 71.01, H: 5.96, N: 13.08 %. Found: C: 70.09, H: 5.98, N: 13.07%.

3. Result and Discussion

The antipyrine based Schiff bases have been synthesized from the condensation reaction of different substituted benzaldehyde with 4-aminoantipyrine under solvent free condition at room temperature with producing excellent yield of products at a short time of duration (Scheme 1) (Table 1).



Scheme 1. Synthesis of 4-Aminoantipyrine Schiff base

Table 1. Characteristic of Synthesized Aminoantipyrine Schiff bases

Compound	Formula	Time/min.	Yield %
C1	C ₁₈ H ₁₇ N ₃ O	40	94
C2	C ₁₈ H ₁₆ FN ₃ O	38	94
C3	C ₁₈ H ₁₆ ClN ₃ O	38	94
C4	C ₁₈ H ₁₅ Cl ₂ N ₃ O	40	95
C5	C ₁₈ H ₁₇ N ₃ O ₂	38	94
C6	C ₂₀ H ₂₁ N ₃ O ₃	39	95
C7	C ₂₄ H ₂₉ N ₃ O ₂	40	93
C8	C ₁₈ H ₁₆ FN ₃ O	39	95
C9	C ₁₈ H ₁₆ N ₄ O ₃	40	92
C10	C ₁₉ H ₁₉ N ₃ O ₂	39	94

Choosing an organic catalyst for producing the desired compounds required several entries of the reaction conditions as well as implementing different type of solvents, then using the DMEA as a catalyst with different concentrations in order to produce and isolate highest amount of the products. The advantageous role of DMEA comes from its bifunctional group (H donor and H acceptor) for the activation of the reaction, the possible interaction of the catalyst is shown in Figure 2.




Fig. 2. Plausible DMEA bifunctional interaction

The yield percent of the model product C1 (benzaldehyde with 4-aminoantipyrene) have been illustrated in Table 2, as well as optimizing the reaction to obtain best amount of the products.

Table 2. Optimization of Reaction Conditions

Entry	Solvent	Catalyst (mol %)	Method	Time/min.	Yield %
1	H ₂ O	–	Reflux	240	83 %
2	EtOH	–	Reflux	130	88 %
3	DMF	–	Reflux	150	89 %
4	–	DMEA (5 mol%)	Stirring	40	93 %
5	–	DMEA (10 mol%)	Stirring	40	93 %
6	–	DMEA (15 mol%)	Stirring	40	93 %
7	H ₂ O	–	U.S.	90	93 %

DMEA= N, N-Dimethylethanolamine, EtOH= Ethanol
U.S.= Ultrasonic bath, DMF= Dimethylformamide

In the presence of water as a solvent (entry 1) the reactants were reacted at reflux temperature for about 4 hours and yielded 83%, but when the solvent has changed to ethanol (entry 2) the yield percent increased to 88% in shorter time nearly 2 hours. In other hand, more time required (3 hours) when dimethyl formamide is used as a solvent but higher amount of the product is obtained (89% yield).

In contrast, in the case of solvent free and room temperature condition, the DMEA has used as a catalyst (entry 3,4, and 5), the reactants were condensed in an excellent amount (93 %). The yields were the same by changing the amount of the catalyst from 5 to 15 mol%, it reveals that, the amount of the catalyst has the same effect on the production of the Schiff bases. On the other hand, the amount of the products synthesized via ultrasonic irradiation (entry 7) has the identical yield with 5 mol % DMEA, but the predominance of the latter comes from its ability to perform the reaction in a shorter time without any solvents at room temperature. The structure of all the synthesized compounds were clarified by FTIR, ¹HNMR, ¹³CNMR, and elemental analysis, and their data are given in the experimental section.

4. Conclusion

In spite of the developments in the production of new and modern chemical compounds for several and variety of applications, still the interest toward the production of Schiff bases is highly considerable due to their importance in several fields and applications, such as biological and pharmaceutical activity, inorganic and coordination chemistry, biochemistry, etc.

Therefore, we have implemented an ecofriendly procedure to synthesize a series of Schiff bases derived from 4-aminoantipyrene and substituted

benzaldehydes by the direct condensation reaction at room temperature under solvent free condition, in which the products were synthesized with excellent yield and simple workout. Choosing the DMEA as an organic catalyst (it is available and inexpensive) in which leads to synthesize the products including two new products ((4-(3,4-dimethoxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one) and (4-(4-hexyloxybenzylideneamino)-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one)) in a very fast reaction method that there is no reports have been documented in this manner, as well as the safety of the catalyst and its easy removal from the product by addition of water only.

Acknowledgements

Financial support by Salahaddin University is gratefully acknowledged.

Dr. Soma Majedi your effort and support are highly appreciated.

References

- [1] S.A. Jarallah, O.A. Nief, A.J.K. Atia, Synthesis, characterization of heterocyclic compounds and preliminary evaluation of their antibacterial activity and antioxidant agents, *Journal of Pharmaceutical Sciences and Research*, 11 (2019) 1010-1015.
- [2] Y. Mi, J. Zhang, X. Han, W. Tan, Q. Miao, J. Cui, Q. Li, Z. Guo, Modification of carboxymethyl inulin with heterocyclic compounds: Synthesis, characterization, antioxidant and antifungal activities, *International Journal of Biological Macromolecules*, 181 (2021) 572-581.
- [3] İ. Şahin, F.B. Özgeriş, M. Köse, E. Bakan, F. Tümer, Synthesis, characterization, and antioxidant and anticancer activity of 1, 4-disubstituted 1, 2, 3-triazoles, *Journal of Molecular Structure*, 1232 (2021) 130042-130050.
- [4] Y. Mi, Q. Li, Q. Miao, W. Tan, J. Zhang, Z. Guo, Enhanced antifungal and antioxidant activities of new chitosan derivatives modified with Schiff base bearing benzenoid/heterocyclic moieties, *International Journal of Biological Macromolecules*, 208 (2022) 586-595.
- [5] R. Kenchappa, Y.D. Bodke, Synthesis, analgesic and anti-inflammatory activity of benzofuran pyrazole heterocycles, *Chemical Data Collections*, 28 (2020) 100453.
- [6] E.A. Fayed, E.M.E. Al-Arab, A.S. Saleh, A.H. Bayoumi, Y.A. Ammar, Design, synthesis, in silico studies, in vivo and in vitro assessment of pyridones and thiazolidinones as anti-inflammatory, antipyretic and ulcerogenic hits, *Journal of Molecular Structure*, 1260 (2022) 132839.
- [7] A. Al-Mulla, A review: biological importance of heterocyclic compounds, *Der Pharma Chemica*, 9 (2017) 141-147.
- [8] O. Mazimba, K. Wale, D. Loeto, T. Kwape, Antioxidant and antimicrobial studies on fused-ring pyrazolones and isoxazolones, *Bioorganic & Medicinal Chemistry*, 22 (2014) 6564-6569.
- [9] N. Uramaru, H. Shigematsu, A. Toda, R. Eyanagi, S. Kitamura, S. Ohta, Design, synthesis, and pharmacological activity of nonallergenic pyrazolone-type antipyretic analgesics, *Journal of medicinal chemistry*, 53 (2010) 8727-8733.

- [10] M. Abdel-Aziz, G.E.-D.A. Abuo-Rahma, A.A. Hassan, Synthesis of novel pyrazole derivatives and evaluation of their antidepressant and anticonvulsant activities, *European journal of medicinal chemistry*, 44 (2009) 3480-3487.
- [11] L. Wang, Z. Yang, T. Ni, W. Shi, Y. Guo, K. Li, A. Shi, S. Wu, C. Sheng, Discovery of pyrazolone spirocyclohexadienone derivatives with potent antitumor activity, *Bioorganic & Medicinal Chemistry Letters*, 30 (2020) 126662-126669.
- [12] F. Marchetti, C. Pettinari, C. Di Nicola, A. Tombesi, R. Pettinari, Coordination chemistry of pyrazolone-based ligands and applications of their metal complexes, *Coordination Chemistry Reviews*, 401 (2019) 213069.
- [13] K.R. Surati, P.A. Sathe, Schiff base pyrazolone complexes of iron (III): synthesis, characterization, antimicrobial and antioxidant activity, *Medicinal chemistry research*, 25 (2016) 2742-2751.
- [14] M.S. Alam, J.-H. Choi, D.-U. Lee, Synthesis of novel Schiff base analogues of 4-amino-1, 5-dimethyl-2-phenylpyrazol-3-one and their evaluation for antioxidant and anti-inflammatory activity, *Bioorganic & medicinal chemistry*, 20 (2012) 4103-4108.
- [15] A. Sakthivel, K. Jeyasubramanian, B. Thangagiri, J.D. Raja, Recent advances in Schiff base metal complexes derived from 4-aminoantipyrine derivatives and their potential applications, *Journal of Molecular Structure*, 1222 (2020) 128885.
- [16] K. Sunand, K.V. Kumar, K. Ashwini, P.S. Kumar, S. Vishnu, A. Samala, Synthesis characterization and antibacterial studies of 4-aminoantipyrine schiff's bases, *International Journal of Applied Pharmaceutical Sciences and Research*, 2 (2017) 8-14.
- [17] H.M. Alabidi, A.M. Farhan, H. Al-labban, A. Aljanaby, New Derivatives from 4-Amino Anti-Pyrine and Vanillin, Synthesis Characterization and Antibacterial Activity, *Egyptian Journal of Chemistry*, (2022).
- [18] R. Çakmak, E. Başaran, M. Boğa, Ö. Erdoğan, E. Çınar, Ö. Çevik, Schiff Base Derivatives of 4-Aminoantipyrine as Promising Molecules: Synthesis, Structural Characterization, and Biological Activities, *Russian Journal of Bioorganic Chemistry*, 48 (2022) 334-344.
- [19] S. Parvarinezhad, M. Salehi, M. Kubicki, Synthesis, characterization, spectral studies and evaluation of noncovalent interactions in co-crystal of μ -oxobridged polymeric copper (II) complex derived from pyrazolone by theoretical studies, *Journal of Molecular Structure*, 1260 (2022) 132780.
- [20] H.M.Y. Al-Labban, H.M. Sadiq, A.A.J. Aljanaby, Synthesis, Characterization and study biological activity of some Schiff bases derivatives from 4-amino antipyrine as a starting material, *Journal of Physics: Conference Series*, IOP Publishing, 2019, 052007-052015.
- [21] M.A. Ashraf, K. Mahmood, A. Wajid, M.J. Maah, I. Yusoff, Synthesis, characterization and biological activity of Schiff bases, *IPCBE*, 10 (2011) 185.
- [22] A.M. Asiri, S.A. Khan, Synthesis and anti-bacterial activities of some novel schiff bases derived from aminophenazone, *molecules*, 15 (2010) 6850-6858.
- [23] C.M. da Silva, M.M. Silva, F.S. Reis, A.L.T. Ruiz, J.E. de Carvalho, J.C. Santos, I.M. Figueiredo, R.B. Alves, L.V. Modolo, Â. de Fátima, Studies on free radical scavenging, cancer cell antiproliferation, and calf thymus DNA interaction of Schiff bases, *Journal of Photochemistry and Photobiology B: Biology*, 172 (2017) 129-138.
- [24] A.A. Ahmed, J.O. Peter, F.A.B. Ngaramu, Synthesis and characterization of mixed ligand metal (II) complexes with Schiff base and 8-hydroxyquinoline as ligands.
- [25] A.A. Ahmed, G.M.G. LEMOS, Synthesis and characterization of Ni (II) complex with Schiff base derived from benzophenone and 2-aminophenol, *Journal of Chemistry Letters*, (2021).
- [26] E. ÇINAR, E. BAŞARAN, Ö. ERDOĞAN, R. ÇAKMAK, B. Mehmet, Ö. ÇEVİK, Synthesis and biological evaluation of some pyrazolone based Schiff base derivatives as enzymes inhibitors, antioxidant, and anticancer agents, (2021).
- [27] I. Kostova, L. Saso, Advances in research of Schiff-base metal complexes as potent antioxidants, *Current medicinal chemistry*, 20 (2013) 4609-4632.
- [28] M.S. Alam, D.-U. Lee, Synthesis, molecular structure and antioxidant activity of (E)-4-[benzylideneamino]-1, 5-dimethyl-2-phenyl-1H-pyrazol-3 (2H)-one, a Schiff base ligand of 4-aminoantipyrine, *Journal of Chemical Crystallography*, 42 (2012) 93-102.
- [29] N. Bashkatova, E. Korotkova, Y.A. Karbainov, A.Y. Yagovkin, A. Bakibaev, Electrochemical, quantum-chemical and antioxidant properties of antipyrine and its derivatives, *Journal of pharmaceutical and biomedical analysis*, 37 (2005) 1143-1147.
- [30] X. Yu, W. Wang, Organocatalysis: asymmetric cascade reactions catalysed by chiral secondary amines, *Organic & biomolecular chemistry*, 6 (2008) 2037-2046.
- [31] A. Shahriza, R. Teimuri-Mofrad, M. Gholamhosseini-Nazari, Synthesis of a new class of Betti bases by the Mannich-type reaction: efficient, facile, solvent-free and one-pot protocol, *Molecular diversity*, 19 (2015) 87-101.